

Predicate logic

Limitations of propositional logic

Propositional logic only allows us to reason about **completed statements** about things, not about the things themselves.

Example

Some chicken cannot fly
All chicken are birds

Some birds cannot fly

this reasoning can not be expressed in propositional logic

Example

Every player except the winner loses a match

Unary predicate (example)


Consider the statement $2m > 3$.

Whether this statement is true or false depends on the value of m (and on the domain of values).

Unary predicate (example)

Consider the statement $2m > 3$.


Whether this statement is true or false depends on the value of m (and on the domain of values).



Unary predicate (example)

Consider the statement $2m > 3$.


Whether this statement is true or false depends on the value of m (and on the domain of values).



Unary predicate (example)

Consider the statement $2m > 3$.

Whether this statement is true or false depends on the value of m (and on the domain of values).



Note: $2m > 3 \stackrel{\text{val}}{=} m > 3/2$ on \mathbb{Z} and \mathbb{R}


$2m > 3 \stackrel{\text{val}}{=} m \geq 2$ on \mathbb{Z} but not on \mathbb{R}

Unary predicate (example)

Consider the statement $2m > 3$.

a unary
relation

Whether this statement is true or false depends on the value of m (and on the domain of values).




Note: $2m > 3 \stackrel{\text{val}}{=} m > 3/2$ on \mathbb{Z} and \mathbb{R}

$2m > 3 \stackrel{\text{val}}{=} m \geq 2$ on \mathbb{Z} but not on \mathbb{R}

Binary predicate (example)

The statement $3m+n > 3$ is a binary predicate on $\mathbb{R} \times \mathbb{N}$.



a binary
relation

Predicates


In general, an n -ary predicate is an n -ary relation.

If it is on a domain D , then it's a relation $P(x_1, \dots, x_n) \subseteq D^n$ or equivalently a function $P: D^n \rightarrow \{0, 1\}$.

Predicates

In general, an n-ary predicate is an n-ary relation.

If it is on a domain D , then it's a relation $P(x_1, \dots, x_n) \subseteq D^n$ or equivalently a function $P: D^n \rightarrow \{0, 1\}$.




true for certain values
of the variables

Predicates

In general, an n-ary predicate is an n-ary relation.

If it is on a domain D , then it's a relation $P(x_1, \dots, x_n) \subseteq D^n$ or equivalently a function $P: D^n \rightarrow \{0, 1\}$.



true for certain values
of the variables

We can turn a predicate, into a proposition in three ways:

1. By assigning values to the variables.
2. By universal quantification.
3. By existential quantification.

Predicates

In general, an n -ary predicate is an n -ary relation.

If it is on a domain D , then it's a relation $P(x_1, \dots, x_n) \subseteq D^n$ or equivalently a function $P: D^n \rightarrow \{0, 1\}$.


$$2m > 3$$



true for certain values
of the variables

We can turn a predicate, into a proposition in three ways:

1. By assigning values to the variables.
2. By universal quantification.
3. By existential quantification.

Predicates

In general, an n-ary predicate is an n-ary relation.

If it is on a domain D , then it's a relation $P(x_1, \dots, x_n) \subseteq D^n$ or equivalently a function $P: D^n \rightarrow \{0, 1\}$.

$$2m > 3$$

true for certain values
of the variables

We can turn a predicate, into a proposition in three ways:

1. By assigning values to the variables.
2. By universal quantification.
3. By existential quantification.

for $m=2$
 $2 \cdot 2 > 3$
is a true proposition

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

For all m in \mathbb{Z} , $2m > 3$

Universal quantification


The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

For all m in \mathbb{Z} , $2m > 3$

false, e.g.
for $m = 1$

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:



false, e.g.
for $m = 1$

For all m in \mathbb{Z} , $2m > 3$

Notation: $\forall_m [m \in \mathbb{Z} : 2m > 3]$

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

false, e.g.
for $m = 1$

For all m in \mathbb{Z} , $2m > 3$

universal
quantifier

Notation:

$\forall m [m \in \mathbb{Z} : 2m > 3]$

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

For all m in \mathbb{Z} , $2m > 3$

false, e.g.
for $m = 1$

universal
quantifier

Notation:

$\forall_m [m \in \mathbb{Z} : 2m > 3]$

domain
(predicate)

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

For all m in \mathbb{Z} , $2m > 3$

false, e.g.
for $m = 1$

Notation:

$\forall_m [m \in \mathbb{Z} : 2m > 3]$

universal
quantifier

domain
(predicate)

predicate

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

false, e.g.
for $m = 1$

For all m in \mathbb{Z} , $2m > 3$

Notation:

$\forall_m [m \in \mathbb{Z} : 2m > 3]$

universal
quantifier

predicate

domain
(predicate)

In general:

$\forall_x [P(x) : Q(x)]$ for “all x satisfying P satisfy Q ”

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

For all m in \mathbb{Z} , $2m > 3$

false, e.g.
for $m = 1$

Notation:

$\forall_m [m \in \mathbb{Z} : 2m > 3]$

universal
quantifier

predicate

domain
(predicate)

other notation:

$\forall x. P(x) \Rightarrow Q(x)$

In general:

$\forall_x [P(x) : Q(x)]$ for “all x satisfying P satisfy Q ”

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

There exists m in \mathbb{Z} , $2m > 3$

Existential quantification


The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

There exists m in \mathbb{Z} , $2m > 3$

true, e.g.
 $m = 2$

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:



true, e.g.
 $m = 2$

There exists m in \mathbb{Z} , $2m > 3$

Notation: $\exists m [m \in \mathbb{Z} : 2m > 3]$

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

true, e.g.
 $m = 2$

There exists m in \mathbb{Z} , $2m > 3$

existential
quantifier

Notation:

$\exists m [m \in \mathbb{Z} : 2m > 3]$

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

true, e.g.
 $m = 2$

There exists m in \mathbb{Z} , $2m > 3$

existential
quantifier

Notation:

$\exists_m [m \in \mathbb{Z} : 2m > 3]$

domain
(predicate)

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

true, e.g.
 $m = 2$

There exists m in \mathbb{Z} , $2m > 3$

Notation:

$\exists_m [m \in \mathbb{Z} : 2m > 3]$

existential
quantifier

domain
(predicate)

predicate

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

true, e.g.
 $m = 2$

There exists m in \mathbb{Z} , $2m > 3$

Notation:

$\exists_m [m \in \mathbb{Z} : 2m > 3]$

existential
quantifier

predicate

domain
(predicate)

In general:

$\exists_x [P(x) : Q(x)]$ for

“there exists x satisfying P that satisfies Q ”

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

true, e.g.
 $m = 2$

There exists m in \mathbb{Z} , $2m > 3$

Notation:

$\exists_m [m \in \mathbb{Z} : 2m > 3]$

existential
quantifier

predicate

domain
(predicate)

other notation:

$\exists x. P(x) \wedge Q(x)$

In general:

$\exists_x [P(x) : Q(x)]$ for

“there exists x satisfying P that satisfies Q ”

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:



in 8
possible
ways

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:



in 8 possible ways

One way is: $\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : 3m + n > 3]]$

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:

in 8 possible ways

One way is: $\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : \underline{3m + n > 3}]]$

binary predicate

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:

in 8 possible ways

One way is:

$$\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : \underline{3m + n > 3}]]$$

unary predicate

binary predicate

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:

in 8 possible ways

One way is:

$$\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : \underline{3m + n > 3}]]$$

unary predicate

binary predicate

proposition,
nullary predicate

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:

in 8 possible ways

One way is:

$$\exists m [m \in \mathbb{R} : \forall n [n \in \mathbb{N} : 3m + n > 3]]$$

unary predicate

binary predicate

proposition,
nullary predicate

other notation:

$$\exists m. (m \in \mathbb{R} \wedge \forall n. (n \in \mathbb{N} \Rightarrow 3m+n > 3))$$

Notation

Notation

We write $\forall_x [P]$ for $\forall_x [T : P]$

Notation

We write $\forall_x [P]$ for $\forall_x [T : P]$



also for \exists

Notation

We write $\forall_x [P]$ for $\forall_x [T : P]$

also for \exists

We also write $\exists_m, \forall_n [(m,n) \in \mathbb{R} \times \mathbb{N} : 3m + n > 3]$

for $\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : 3m + n > 3]]$

Notation

We write $\forall_x [P]$ for $\forall_x [T : P]$

also for \exists

We also write $\exists_m, \forall_n [(m,n) \in \mathbb{R} \times \mathbb{N} : 3m + n > 3]$

for $\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : 3m + n > 3]]$

And even $\exists_{m,n} [(m,n) \in \mathbb{R} \times \mathbb{N} : 3m + n > 3]$

for $\exists_m [m \in \mathbb{R} : \exists_n [n \in \mathbb{N} : 3m + n > 3]]$

Notation

We write $\forall_x [P]$ for $\forall_x [T : P]$

also for \exists

We also write $\exists_m, \forall_n [(m,n) \in \mathbb{R} \times \mathbb{N} : 3m + n > 3]$

for $\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : 3m + n > 3]]$

And even $\exists_{m,n} [(m,n) \in \mathbb{R} \times \mathbb{N} : 3m + n > 3]$

for $\exists_m [m \in \mathbb{R} : \exists_n [n \in \mathbb{N} : 3m + n > 3]]$

but only for the same
quantifier!

Quantification - task

Let P be the set of all tennis players.

Let $w \in P$ be the winner.

For $p, q \in P$, write $p \neq q$ for “ p and q are different players” .

Let M be the set of all matches.

For $p \in P$ and $m \in M$, write $L(p,m)$ for

“player p loses match m ”.

Write the following sentence as a formula with predicates and quantifiers:

Every player except the winner loses a match.

Quantification - task

Let P be the set of all tennis players.

Let $w \in P$ be the winner.

Thanks to Bas Luttik

For $p, q \in P$, write $p \neq q$ for “ p and q are different players” .

Let M be the set of all matches.

For $p \in P$ and $m \in M$, write $L(p,m)$ for

“player p loses match m ”.

Write the following sentence as a formula with predicates and quantifiers:

Every player except the winner loses a match.

Equivalences with quantifiers

Renaming bound variables

Bound variables

$$\forall_x [P:Q] \stackrel{val}{=} \forall_y [P[y/x]:Q[y/x]]$$

$$\exists_x [P:Q] \stackrel{val}{=} \exists_y [P[y/x]:Q[y/x]]$$

if y does not occur in
 P or Q (not even in $\forall y, \exists y$)

Domain splitting

Examples:

$$\begin{aligned} & \forall x [x \leq 1 \vee x \geq 5 : x^2 - 6x + 5 \geq 0] \\ \stackrel{val}{=} & \forall x [x \leq 1 : x^2 - 6x + 5 \geq 0] \wedge \forall x [x \geq 5 : x^2 - 6x + 5 \geq 0] \end{aligned}$$

Domain splitting

Examples:

$$\begin{aligned} & \forall x [x \leq 1 \vee x \geq 5 : x^2 - 6x + 5 \geq 0] \\ \stackrel{val}{=} & \forall x [x \leq 1 : x^2 - 6x + 5 \geq 0] \wedge \forall x [x \geq 5 : x^2 - 6x + 5 \geq 0] \end{aligned}$$

$$\begin{aligned} & \exists k [0 \leq k \leq n : k^2 \leq 10] \\ \stackrel{val}{=} & \exists k [0 \leq k \leq n - 1 \vee k = n : k^2 \leq 10] \\ \stackrel{val}{=} & \exists k [0 \leq k \leq n - 1 : k^2 \leq 10] \vee \exists k [k = n : k^2 \leq 10] \end{aligned}$$

Domain splitting

Examples:

$$\begin{aligned} & \forall x [x \leq 1 \vee x \geq 5 : x^2 - 6x + 5 \geq 0] \\ \stackrel{val}{=} & \forall x [x \leq 1 : x^2 - 6x + 5 \geq 0] \wedge \forall x [x \geq 5 : x^2 - 6x + 5 \geq 0] \end{aligned}$$

$$\begin{aligned} & \exists k [0 \leq k \leq n : k^2 \leq 10] \\ \stackrel{val}{=} & \exists k [0 \leq k \leq n - 1 \vee k = n : k^2 \leq 10] \\ \stackrel{val}{=} & \exists k [0 \leq k \leq n - 1 : k^2 \leq 10] \vee \exists k [k = n : k^2 \leq 10] \end{aligned}$$

Domain splitting

$$\forall x [P \vee Q : R] \stackrel{val}{=} \forall x [P : R] \wedge \forall x [Q : R]$$

$$\exists x [P \vee Q : R] \stackrel{val}{=} \exists x [P : R] \vee \exists x [Q : R]$$

Equivalences with quantifiers

One-element domain

$$\forall x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

$$\exists x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

Equivalences with quantifiers

One-element domain

$$\forall x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

$$\exists x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

Example:

$$\forall x [x = 3 : 2 \cdot x \geq 1] \stackrel{val}{=} 2 \cdot 3 \geq 1$$

Equivalences with quantifiers

One-element domain

$$\forall x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

$$\exists x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

Example:

$$\forall x [x = 3 : 2 \cdot x \geq 1] \stackrel{val}{=} 2 \cdot 3 \geq 1$$

Empty domain

$$\forall x [F : Q] \stackrel{val}{=} T$$

$$\exists x [F : Q] \stackrel{val}{=} F$$

Equivalences with quantifiers

One-element domain

$$\forall x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

$$\exists x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

Example:

$$\forall x [x = 3 : 2 \cdot x \geq 1] \stackrel{val}{=} 2 \cdot 3 \geq 1$$

“All Marsians are green”

Empty domain

$$\forall x [F : Q] \stackrel{val}{=} T$$

$$\exists x [F : Q] \stackrel{val}{=} F$$

Domain weakening

Intuition: The following are equivalent

$$\forall x [x \in D : A(x)] \quad \text{and} \quad \forall x [x \in D \Rightarrow A(x)]$$

$$\exists x [x \in D : A(x)] \quad \text{and} \quad \exists x [x \in D \wedge A(x)]$$

The same can be done to parts of the domain

Domain weakening

Intuition: The following are equivalent

$$\begin{array}{l} \forall x [x \in D : A(x)] \quad \text{and} \quad \forall x [x \in D \Rightarrow A(x)] \\ \exists x [x \in D : A(x)] \quad \text{and} \quad \exists x [x \in D \wedge A(x)] \end{array}$$

The same can be done to parts of the domain

Domain weakening

$$\forall x [P \wedge Q : R] \stackrel{val}{=} \forall x [P : Q \Rightarrow R]$$

$$\exists x [P \wedge Q : R] \stackrel{val}{=} \exists x [P : Q \wedge R]$$

Domain weakening

Intuition: The following are equivalent

$$\begin{array}{l} \forall x [x \in D : A(x)] \quad \text{and} \quad \forall x [x \in D \Rightarrow A(x)] \\ \exists x [x \in D : A(x)] \quad \text{and} \quad \exists x [x \in D \wedge A(x)] \end{array}$$

The same can be done to parts of the domain

Domain weakening

$$\begin{array}{l} \forall x [P \wedge Q : R] \stackrel{val}{=} \forall x [P : Q \Rightarrow R] \\ \exists x [P \wedge Q : R] \stackrel{val}{=} \exists x [P : Q \wedge R] \end{array}$$

$$P \wedge Q \stackrel{val}{\models} P$$

De Morgan with quantifiers

De Morgan

$$\neg \forall x [P : Q] \stackrel{val}{=} \exists x [P : \neg Q]$$

$$\neg \exists x [P : Q] \stackrel{val}{=} \forall x [P : \neg Q]$$

De Morgan with quantifiers

De Morgan

$$\neg \forall x [P : Q] \stackrel{val}{=} \exists x [P : \neg Q]$$
$$\neg \exists x [P : Q] \stackrel{val}{=} \forall x [P : \neg Q]$$

not for all = at least for one not

not exists = for all not

De Morgan with quantifiers

De Morgan

$$\neg \forall x [P : Q] \stackrel{val}{=} \exists x [P : \neg Q]$$
$$\neg \exists x [P : Q] \stackrel{val}{=} \forall x [P : \neg Q]$$

not for all = at least for one not

not exists = for all not

Hence: $\neg \forall = \exists \neg$ and $\neg \exists = \forall \neg$

De Morgan with quantifiers

De Morgan

$$\neg \forall x [P : Q] \stackrel{val}{=} \exists x [P : \neg Q]$$

$$\neg \exists x [P : Q] \stackrel{val}{=} \forall x [P : \neg Q]$$

not for all = at least for one not

not exists = for all not

Hence: $\neg \forall = \exists \neg$ and $\neg \exists = \forall \neg$

It holds further that:

$$\neg \forall x \neg = \exists x \neg \neg = \exists x$$

$$\neg \exists x \neg = \forall x \neg \neg = \forall x$$

Substitution

meta rule

Simple

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P] \stackrel{val}{=} \psi[\xi/P]}$$

Sequential

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P][\eta/Q] \stackrel{val}{=} \psi[\xi/P][\eta/Q]}$$

Simultaneous

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P, \eta/Q] \stackrel{val}{=} \psi[\xi/P, \eta/Q]}$$

EVERY occurrence of P is substituted!

holds also for
quantified formulas!

Substitution

meta rule

Simple

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P] \stackrel{val}{=} \psi[\xi/P]}$$

Sequential

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P][\eta/Q] \stackrel{val}{=} \psi[\xi/P][\eta/Q]}$$

Simultaneous

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P, \eta/Q] \stackrel{val}{=} \psi[\xi/P, \eta/Q]}$$

EVERY occurrence of
P is substituted!

The rule of Leibniz

Leibniz

$$\phi \stackrel{val}{=} \psi$$

$$C[\phi] \stackrel{val}{=} C[\psi]$$

meta rule

formula that has
 ϕ as a sub formula

single occurrence is
replaced!

holds also for
quantified formulas!

The rule of Leibnitz

meta rule

Leibnitz

$$\phi \stackrel{val}{=} \psi$$

$$C[\phi] \stackrel{val}{=} C[\psi]$$

formula that has
 ϕ as a sub formula

single occurrence is
replaced!

Other equivalences with quantifiers

Exchange trick

$$\forall x [P:Q] \stackrel{val}{=} \forall x [\neg Q:\neg P]$$

$$\exists x [P:Q] \stackrel{val}{=} \exists x [Q:P]$$

Other equivalences with quantifiers

Exchange trick

$$\forall x [P:Q] \stackrel{val}{=} \forall x [\neg Q:\neg P]$$

$$\exists x [P:Q] \stackrel{val}{=} \exists x [Q:P]$$

No wonder as

$$\forall x [P:Q] \stackrel{val}{=} \forall x [P \Rightarrow Q]$$

$$\exists x [P:Q] \stackrel{val}{=} \exists x [P \wedge Q]$$

Other equivalences with quantifiers

Exchange trick

$$\forall_x [P:Q] \stackrel{val}{=} \forall_x [\neg Q:\neg P]$$

$$\exists_x [P:Q] \stackrel{val}{=} \exists_x [Q:P]$$

No wonder as

$$\forall_x [P:Q] \stackrel{val}{=} \forall_x [P \Rightarrow Q]$$

$$\exists_x [P:Q] \stackrel{val}{=} \exists_x [P \wedge Q]$$

Term splitting

$$\forall_x [P:Q \wedge R] \stackrel{val}{=} \forall_x [P:Q] \wedge \forall_x [P:R]$$

$$\exists_x [P:Q \vee R] \stackrel{val}{=} \exists_x [P:Q] \vee \exists_x [P:R]$$

Other equivalences with quantifiers

Monotonicity of quantifiers

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\forall x [P:Q] \Rightarrow \forall x [P:R]) \stackrel{val}{=} T$$

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\exists x [P:Q] \Rightarrow \exists x [P:R]) \stackrel{val}{=} T$$

Other equivalences with quantifiers

Monotonicity of quantifiers

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\forall x [P:Q] \Rightarrow \forall x [P:R]) \stackrel{val}{=} T$$

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\exists x [P:Q] \Rightarrow \exists x [P:R]) \stackrel{val}{=} T$$

tautologies

Other equivalences with quantifiers

Monotonicity of quantifiers

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\forall x [P:Q] \Rightarrow \forall x [P:R]) \stackrel{val}{=} T$$

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\exists x [P:Q] \Rightarrow \exists x [P:R]) \stackrel{val}{=} T$$

tautologies

Lemma E1: $P \stackrel{val}{=} Q$ iff $P \Leftrightarrow Q$ is a tautology.

Other equivalences with quantifiers

Monotonicity of quantifiers

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\forall x [P:Q] \Rightarrow \forall x [P:R]) \stackrel{val}{=} T$$

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\exists x [P:Q] \Rightarrow \exists x [P:R]) \stackrel{val}{=} T$$

tautologies

Lemma E1: $P \stackrel{val}{=} Q$ iff $P \Leftrightarrow Q$ is a tautology.

Lemma W4: $P \stackrel{val}{\models} Q$ iff $P \Rightarrow Q$ is a tautology.

Other equivalences with quantifiers

Monotonicity of quantifiers

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\forall x [P:Q] \Rightarrow \forall x [P:R]) \stackrel{val}{=} T$$

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\exists x [P:Q] \Rightarrow \exists x [P:R]) \stackrel{val}{=} T$$

tautologies

Lemma E1: $P \stackrel{val}{=} Q$ iff $P \Leftrightarrow Q$ is a tautology.

Lemma W4: $P \stackrel{val}{\models} Q$ iff $P \Rightarrow Q$ is a tautology.

still hold (in predicate logic)

Other equivalences with quantifiers

Monotonicity of quantifiers

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\forall x [P:Q] \Rightarrow \forall x [P:R]) \stackrel{val}{=} T$$

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\exists x [P:Q] \Rightarrow \exists x [P:R]) \stackrel{val}{=} T$$

tautologies

Lemma E1: $P \stackrel{val}{=} Q$ iff $P \Leftrightarrow Q$ is a tautology.

Lemma W4: $P \stackrel{val}{\models} Q$ iff $P \Rightarrow Q$ is a tautology.

Lemma W5: If $Q \stackrel{val}{\models} R$ then $\forall x [P:Q] \stackrel{val}{\models} \forall x [P:R]$.

still hold (in predicate logic)