


Functions, mappings


Def. If A and B are sets, a function (mapping, *Abbildung*) f from A to B , notation $f: A \longrightarrow B$ is an assignment (of elements of B to elements of A , we write $f(a)$ for the element assigned to a) s. t.
for every $a \in A$, there exists a unique $b \in B$ such that $b = f(a)$.



$\{(a, f(a)) \mid a \in A\}$ is the *graph* of the function f

Functions, mappings

When $f: A \longrightarrow B$ then $\text{dom } f = A$ and $\text{cod } f = B$



Let $f: A \longrightarrow B$ and $A' \subseteq A$.

The image (**Bild**) of A' is the set $f(A') = \{f(a) \mid a \in A'\} \subseteq B$.

$$f(A') = \{b \in B \mid \text{there is an } a \in A' \text{ with } b = f(a)\}$$

if $a \in A'$, then $f(a) \in f(A')$

So f extends to a function $f: \mathcal{P}(A) \longrightarrow \mathcal{P}(B)$, the image-function.

Functions, mappings

Let $f: A \longrightarrow B$ and $B' \subseteq B$.

The inverse image (**Urbild**) of B' is the set

$$f^{-1}(B') = \{a \mid f(a) \in B'\} \subseteq A.$$


$$a \in f^{-1}(B') \quad \text{iff} \quad f(a) \in B'$$

Again the inverse image induces a function $f^{-1}: \mathcal{P}(B) \longrightarrow \mathcal{P}(A)$, the inverse-image-function.

Lemma F1: Let $f: A \longrightarrow B$, $A' \subseteq A$, and $B' \subseteq B$. Then

$$A' \subseteq f^{-1}(f(A')) \quad \text{and} \quad f(f^{-1}(B')) \subseteq B'$$

(in general no more than this holds)

Equality of functions

Let $f:A \longrightarrow B$ and $g:C \longrightarrow D$

Def. The functions $f:A \longrightarrow B$ and $g:C \longrightarrow D$ are equal iff

(1) $A = C$

(2) $B = D$

(3) for all $a \in A$, $f(a) = g(a)$.





$\text{dom } f = \text{dom } g$

$\text{cod } f = \text{cod } g$

Recall...


Def. If A and B are sets, a function f from A to B , notation $f: A \longrightarrow B$ is an assignment s. t.

for every $a \in A$, there exists a unique $b \in B$ such that $b = f(a)$.




Special functions

The number of ingoing arrows for a function can be 0, 1, or more. Based on this, we distinguish some special functions.



at most one incoming arrow
injection




at least one incoming arrow
surjection


exactly one incoming arrow (injection + surjection) bijection

Special functions

Def. A function $f:A \longrightarrow B$ is injective iff
for all $a, b \in A$, if $f(a) = f(b)$ then $a = b$.



Def. A function $f:A \longrightarrow B$ is surjective iff
for all $b \in B$, there exists $a \in A$ such that $f(a) = b$.



Def. A function $f:A \longrightarrow B$ is bijective iff
 f is injective and surjective.

Simple characterisations

Lemma I: A function $f:A \longrightarrow B$ is injective iff
for all $b \in B$, $|f^{-1}(\{b\})| \leq 1$.

at most one incoming arrow
injection

Lemma S: A function $f:A \longrightarrow B$ is surjective iff
 $|f^{-1}(\{b\})| \geq 1$ for all $b \in B$ iff
 $f(A) = B$.

at least one incoming arrow
surjection

Lemma B: A function $f:A \longrightarrow B$ is bijective iff
 $|f^{-1}(\{b\})| = 1$ for all $b \in B$ iff
 f is both injective and surjective.

exactly one incoming arrow
bijection

Some properties

Lemma I2: Let $f:A \longrightarrow B$ be injective and let $A' \subseteq A$. Then
 $f(x) \in f(A')$ iff $x \in A'$.


if holds always!

Prop. I3: Let $f:A \longrightarrow B$ be injective and let $A' \subseteq A$. Then
 $f^{-1}(f(A')) = A'$.

Prop. S2: Let $f:A \longrightarrow B$ be surjective and let $B' \subseteq B$. Then
 $f(f^{-1}(B')) = B'$.

Inverse function

Let $f:A \longrightarrow B$ be a **bijection**



well defined only if f is bijective!

Def. The inverse function $f^{-1}: B \longrightarrow A$ is defined as

$$f^{-1}(b) = a \text{ iff } f(a) = b, \quad b \in B.$$

Lemma B2: The inverse function f^{-1} for a bijection f is bijective.

Function composition

Let $f:A \longrightarrow B$ and $g:B \longrightarrow C$

Function composition

Let $f:A \longrightarrow B$ and $g:B \longrightarrow C$

“after”

$g \circ f : A \longrightarrow B \longrightarrow C$

Def. The composition $g \circ f$ is a function $g \circ f : A \longrightarrow C$ given by
 $g \circ f (a) = g(f(a))$, for $a \in A$.


Lemma I4: Let $f:A \longrightarrow B$ and $g:B \longrightarrow C$ be injective. Then
 $g \circ f$ is injective.

Lemma S3: Let $f:A \longrightarrow B$ and $g:B \longrightarrow C$ be surjective. Then
 $g \circ f$ is surjective.

Corollary B2: Let $f:A \longrightarrow B$ and $g:B \longrightarrow C$ be bijective. Then so is $g \circ f$.

A characterization of bijections

Theorem B3: A function $f:A \longrightarrow B$ is bijective iff there exists a function $g:B \longrightarrow A$ with $g \circ f = \text{id}_A$ and $f \circ g = \text{id}_B$.


$$\begin{aligned} \text{id}_A: A &\longrightarrow A, \\ \text{id}_A(a) &= a, \text{ for all } a \in A \end{aligned}$$