

Predicate logic

Limitations of propositional logic

Propositional logic only allows us to reason about **completed statements** about things, not about the things themselves.

Example

Some chicken cannot fly
All chicken are birds

Some birds cannot fly

this reasoning can not
be expressed in
propositional logic

Example


Every player except the winner loses a match

Unary predicate (example)

Consider the statement $2m > 3$.

a unary
relation

Whether this statement is true or false depends on the value of m (and on the domain of values).




Note: $2m > 3 \stackrel{\text{val}}{=} m > 3/2$ on \mathbb{Z} and \mathbb{R}

$2m > 3 \stackrel{\text{val}}{=} m \geq 2$ on \mathbb{Z} but not on \mathbb{R}

Binary predicate (example)

The statement $3m+n > 3$ is a binary predicate on $\mathbb{R} \times \mathbb{N}$.



a binary
relation

Predicates

In general, an n-ary predicate is an n-ary relation.

If it is on a domain D , then it's a relation $P(x_1, \dots, x_n) \subseteq D^n$ or equivalently a function $P: D^n \rightarrow \{0, 1\}$.

$$2m > 3$$

true for certain values
of the variables

We can turn a predicate, into a proposition in three ways:

1. By assigning values to the variables.
2. By universal quantification.
3. By existential quantification.

for $m=2$
 $2 \cdot 2 > 3$
is a true proposition

Universal quantification

The unary predicate $2m > 3$ on \mathbb{Z} can be turned into a proposition by universal quantification:

For all m in \mathbb{Z} , $2m > 3$

false, e.g.
for $m = 1$

Notation:

$\forall_m [m \in \mathbb{Z} : 2m > 3]$

universal
quantifier

domain
(predicate)

predicate

standard (!)
notation:

$\forall x (P(x) \Rightarrow Q(x))$

$\forall x. P(x) \Rightarrow Q(x)$

In general:

$\forall_x [P(x) : Q(x)]$ for “all x satisfying P satisfy Q ”

Existential quantification

The unary predicate $2m > 3$ on \mathbb{Z} can also be turned into a proposition by existential quantification:

There exists m in \mathbb{Z} , $2m > 3$

true, e.g.
 $m = 2$

Notation:

$\exists_m [m \in \mathbb{Z} : 2m > 3]$

existential
quantifier

domain
(predicate)

predicate

standard (!)
notation:

$\exists x (P(x) \wedge Q(x))$

$\exists x. P(x) \wedge Q(x)$

In general:

$\exists_x [P(x) : Q(x)]$ for

“there exists x satisfying P that satisfies Q ”

Quantification

The binary predicate $3m+n > 3$ on $\mathbb{R} \times \mathbb{N}$ can also be turned into a proposition by quantification:

in 8 possible ways

One way is:

$$\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : 3m + n > 3]]$$

standard (!) notation:

$$\exists m (m \in \mathbb{R} \wedge \forall n (n \in \mathbb{N} \Rightarrow 3m+n > 3))$$

unary predicate

binary predicate

proposition,
nullary predicate

Notation

We write $\forall_x [P]$ for $\forall_x [T : P]$

also for \exists

We also write $\exists_m, \forall_n [(m,n) \in \mathbb{R} \times \mathbb{N} : 3m + n > 3]$

for $\exists_m [m \in \mathbb{R} : \forall_n [n \in \mathbb{N} : 3m + n > 3]]$

And even $\exists_{m,n} [(m,n) \in \mathbb{R} \times \mathbb{N} : 3m + n > 3]$

for $\exists_m [m \in \mathbb{R} : \exists_n [n \in \mathbb{N} : 3m + n > 3]]$

but only for the same
quantifier!

Quantification - task

Let P be the set of all tennis players.

Let $w \in P$ be the winner.

Thanks to Bas Luttik

For $p, q \in P$, write $p \neq q$ for “ p and q are different players”.

Let M be the set of all matches.

For $p \in P$ and $m \in M$, write $L(p,m)$ for

“player p loses match m ”.

Write the following sentence as a formula with predicates and quantifiers:

Every player except the winner loses a match.

Equivalences with quantifiers

Renaming bound variables

Bound variables

$$\forall_x [P:Q] \stackrel{val}{=} \forall_y [P[y/x]:Q[y/x]]$$

$$\exists_x [P:Q] \stackrel{val}{=} \exists_y [P[y/x]:Q[y/x]]$$

if y does not occur in
P or Q (not even in $\forall y, \exists y$)

Domain splitting

Examples:

$$\begin{aligned} & \forall x [x \leq 1 \vee x \geq 5 : x^2 - 6x + 5 \geq 0] \\ \stackrel{val}{=} & \forall x [x \leq 1 : x^2 - 6x + 5 \geq 0] \wedge \forall x [x \geq 5 : x^2 - 6x + 5 \geq 0] \end{aligned}$$

$$\begin{aligned} & \exists k [0 \leq k \leq n : k^2 \leq 10] \\ \stackrel{val}{=} & \exists k [0 \leq k \leq n - 1 \vee k = n : k^2 \leq 10] \\ \stackrel{val}{=} & \exists k [0 \leq k \leq n - 1 : k^2 \leq 10] \vee \exists k [k = n : k^2 \leq 10] \end{aligned}$$

Domain splitting

$$\forall x [P \vee Q : R] \stackrel{val}{=} \forall x [P : R] \wedge \forall x [Q : R]$$

$$\exists x [P \vee Q : R] \stackrel{val}{=} \exists x [P : R] \vee \exists x [Q : R]$$

Equivalences with quantifiers

One-element domain

$$\forall x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

$$\exists x [x = n : Q] \stackrel{val}{=} Q[n/x]$$

Example:

$$\forall x [x = 3 : 2 \cdot x \geq 1] \stackrel{val}{=} 2 \cdot 3 \geq 1$$

“All Marsians are green”

Empty domain

$$\forall x [F : Q] \stackrel{val}{=} T$$

$$\exists x [F : Q] \stackrel{val}{=} F$$

Domain weakening

Intuition: The following are equivalent

$$\begin{array}{l} \forall x [x \in D : A(x)] \quad \text{and} \quad \forall x [x \in D \Rightarrow A(x)] \\ \exists x [x \in D : A(x)] \quad \text{and} \quad \exists x [x \in D \wedge A(x)] \end{array}$$

The same can be done to parts of the domain

Domain weakening

$$\forall x [P \wedge Q : R] \stackrel{val}{=} \forall x [P : Q \Rightarrow R]$$

$$\exists x [P \wedge Q : R] \stackrel{val}{=} \exists x [P : Q \wedge R]$$

$$P \wedge Q \stackrel{val}{\models} P$$

De Morgan with quantifiers

De Morgan

$$\neg \forall x [P : Q] \stackrel{val}{=} \exists x [P : \neg Q]$$

$$\neg \exists x [P : Q] \stackrel{val}{=} \forall x [P : \neg Q]$$

not for all = at least for one not

not exists = for all not

Hence: $\neg \forall = \exists \neg$ and $\neg \exists = \forall \neg$

It holds further that:

$$\neg \forall x \neg = \exists x \neg \neg = \exists x$$

$$\neg \exists x \neg = \forall x \neg \neg = \forall x$$

holds also for
quantified formulas!

Substitution

meta rule

Simple

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P] \stackrel{val}{=} \psi[\xi/P]}$$

Sequential

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P][\eta/Q] \stackrel{val}{=} \psi[\xi/P][\eta/Q]}$$

Simultaneous

$$\frac{\phi \stackrel{val}{=} \psi}{\phi[\xi/P, \eta/Q] \stackrel{val}{=} \psi[\xi/P, \eta/Q]}$$

EVERY occurrence of
P is substituted!

holds also for
quantified formulas!

The rule of Leibniz

meta rule

Leibniz

$$\phi \stackrel{val}{=} \psi$$

$$C[\phi] \stackrel{val}{=} C[\psi]$$

formula that has
 ϕ as a sub formula

single occurrence is
replaced!

Other equivalences with quantifiers

Exchange trick

$$\forall x [P:Q] \stackrel{val}{=} \forall x [\neg Q:\neg P]$$

$$\exists x [P:Q] \stackrel{val}{=} \exists x [Q:P]$$

No wonder as

$$\forall x [P:Q] \stackrel{val}{=} \forall x [P \Rightarrow Q]$$

$$\exists x [P:Q] \stackrel{val}{=} \exists x [P \wedge Q]$$

Term splitting

$$\forall x [P:Q \wedge R] \stackrel{val}{=} \forall x [P:Q] \wedge \forall x [P:R]$$

$$\exists x [P:Q \vee R] \stackrel{val}{=} \exists x [P:Q] \vee \exists x [P:R]$$

Other equivalences with quantifiers

Monotonicity of quantifiers

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\forall x [P:Q] \Rightarrow \forall x [P:R]) \stackrel{val}{=} T$$

$$\forall x [P:Q \Rightarrow R] \Rightarrow (\exists x [P:Q] \Rightarrow \exists x [P:R]) \stackrel{val}{=} T$$

tautologies

Lemma E1: $P \stackrel{val}{=} Q$ iff $P \Leftrightarrow Q$ is a tautology.

Lemma W4: $P \stackrel{val}{\models} Q$ iff $P \Rightarrow Q$ is a tautology.

Lemma W5: If $Q \stackrel{val}{\models} R$ then $\forall x [P:Q] \stackrel{val}{\models} \forall x [P:R]$.

still hold (in predicate logic)