

# Sets

- A **set**  $S$  is a collection of different objects, the elements of  $S$
- We write  $x \in S$  for ‘ $x$  is an element of  $S$ ’
- A set ‘can’ be specified by
  - (1) listing its elements, e.g.  $S = \{1, 3, 7, 18\}$
  - (2) specifying a **property**, e.g.  $S = \{x \mid P(x)\}$
- Sets can be **finite** e.g.  $\{\diamondsuit, \heartsuit\}$  or **infinite** e.g.  $\mathbb{N}$
- The set with no elements is the **empty set**, notation  $\emptyset$
- The ‘number’ of elements in a set  $S$  is the **cardinality** of  $S$ , notation  $|S|$


P is a proposition over  $x$ , which is true or false

# Sets - properties

- All elements of a set are different
- The elements of a set are not ordered
- The same set can be specified in different ways, e.g.  
 $\{1,2,3,4\}, \{2,3,1,4\}, \{i \mid i \in \mathbb{N} \text{ and } 0 < i < 5\}$

# Subsets, equality

Def.  $A \subseteq B$  iff all elements of A are elements of B  
[iff for all  $a$ , if  $a \in A$ , then  $a \in B$ ]




Def.  $A = B$  iff  $A \subseteq B$  and  $B \subseteq A$

Def.  $A \subset B$  iff  $A \subseteq B$  and  $A \neq B$


# Operations on sets

Def. Union (Vereinigung)  $A \cup B = \{x \mid x \in A \text{ or } x \in B\}$




Def. Intersection (Durchschnitt)  $A \cap B = \{x \mid x \in A \text{ and } x \in B\}$

A and B are disjoint if  $A \cap B = \emptyset$



Def. Difference (Differenz)  $A \setminus B = \{x \mid x \in A \text{ and } x \notin B\}$



Def. Direct product (Kartesisches Produkt)

$(A \times B) \times C \neq A \times (B \times C)$

$A \times B = \{(x,y) \mid x \in A \text{ and } y \in B\}$

ordered pairs

Def. Powerset (Potenzmenge)  $\mathcal{P}(A) = \{X \mid X \subseteq A\}$

# Russell's paradox


- Let  $P$  be the set of all sets that are not an element of itself
- Hence,  $P = \{ x \mid x \notin x\}$
- Is  $P \in P$  ?
- **Contradiction!**

The need for a universal set  $U$

$$S = \{x \mid x \in U \text{ and } P(x)\}$$

# Operations on sets


Def. Difference (Differenz)  $A \setminus B = \{x \mid x \in A \text{ and } x \notin B\}$



Given a universal set  $U$

Def. Complement (Komplement)  $A^c = \{x \mid x \in U \text{ and } x \notin A\}$

Hence  $A^c = U \setminus A$



# Properties of sets

1.  $\emptyset \subseteq X$
2. If  $A \subseteq B$  and  $B \subseteq C$ , then  $A \subseteq C$
3.  $X \cap Y \subseteq X$ ,  $X \cap Y \subseteq Y$
4.  $X \subseteq X \cup Y$ ,  $Y \subseteq X \cup Y$
5. If  $X' \subseteq Y'$  and  $X'' \subseteq Y''$ , then  $X' \cap X'' \subseteq Y' \cap Y''$
6. If  $X' \subseteq Y'$  and  $X'' \subseteq Y''$ , then  $X' \cup X'' \subseteq Y' \cup Y''$
7.  $X \cap Y = X$  iff  $X \subseteq Y$
8.  $X \cap X = X$  (idempotence)
9.  $X \cup X = X$  (idempotence)
10.  $X \cap \emptyset = \emptyset$

# Properties of sets

I1.  $X \cup \emptyset = X$

I2.  $X \cap Y = Y \cap X$  (commutativity)

I3.  $X \cup Y = Y \cup X$  (commutativity)

I4.  $X \cap (Y \cap Z) = (X \cap Y) \cap Z$  (associativity)

I5.  $X \cup (Y \cup Z) = (X \cup Y) \cup Z$  (associativity)

I6.  $X \cap (X \cup Y) = X$  (absorption)

I7.  $X \cup (X \cap Y) = X$  (absorption)

I8.  $X \cap (Y \cup Z) = (X \cap Y) \cup (X \cap Z)$  (distributivity)

I9.  $X \cup (Y \cap Z) = (X \cup Y) \cap (X \cup Z)$  (distributivity)

I10.  $X \setminus Y \subseteq X$

# Properties of sets

$$21. (X \setminus Y) \cap Y = \emptyset$$

$$22. X \cup Y = X \cup (Y \setminus X)$$

$$23. X \setminus X = \emptyset$$

$$24. X \setminus \emptyset = X$$

$$25. \emptyset \setminus X = \emptyset$$

$$26. \text{ If } X \subseteq Y, \text{ then } X \setminus Y = \emptyset$$

$$27. (X^c)^c = X$$

$$28. (X \cap Y)^c = X^c \cup Y^c \quad (\text{De Morgan})$$

$$29. (X \cup Y)^c = X^c \cap Y^c \quad (\text{De Morgan})$$

$$30. X \times \emptyset = \emptyset \quad \emptyset \times X = \emptyset$$

$$31. \emptyset \times X = \emptyset$$

$$32. \text{ If } X \subseteq Y, \text{ then } \mathcal{P}(X) \subseteq \mathcal{P}(Y)$$