

Formale Systeme Proseminar

Tasks for Week 11

Task 1 Rewrite each of the following formulas with a calculation to a simpler formula:

- (a) $\forall x[P : T]$,
- (b) $\forall x[P : F]$
- (c) $\exists x[P : T]$
- (d) $\exists x[P : F]$

Task 2 Rewrite each of the following formulas with a calculation to a simpler formula:

- (a) $\forall x[P \vee Q : \neg P]$,
- (b) $\forall x[P \wedge Q : \neg P]$,
- (c) $\exists x[P \vee Q : \neg P]$,
- (d) $\exists x[P \wedge Q : \neg P]$

Task 3 Show with a calculation that

- (a) $\exists x[P : Q] \stackrel{val}{=} \neg \forall x[Q : \neg P]$,
- (b) $\forall x[P : Q \vee R] \stackrel{val}{=} \forall x[P \wedge \neg Q : R]$.

Task 4 Show with derivations that the following formula is a tautology

$$((P \Rightarrow Q) \Rightarrow P) \Rightarrow ((P \Rightarrow Q) \Rightarrow Q)$$

Task 5 For each of the line numbers of your solution to Task 4, say where the proposition which occurs on that line is valid (i.e. allowed to be used).

Task 6 Show with derivations that the following formula is a tautology

$$\neg(P \Rightarrow Q) \Rightarrow \neg Q$$

Task 7 Give logical derivation of the following tautology

$$(P \wedge \neg Q) \Rightarrow \neg(P \Rightarrow Q)$$

Task 8 Give logical derivation of the following tautology

$$(\neg P \Rightarrow P) \Rightarrow P$$