

Induction

$$P(0) \wedge \forall i [i \in \mathbb{N} : P(i) \Rightarrow P(i+1)] \Rightarrow \forall n [n \in \mathbb{N} : P(n)]$$

P - unary predicate
over \mathbb{N}

...

(m) P(0)
{Assume}

(k) **var** i; i \in \mathbb{N}

(k+1) P(i)
...

(l-1) P(i+1)
{ \Rightarrow -intro on (k+1) and (l-1)}

(l) P(i) \Rightarrow P(i+1)
{ \forall -intro on (k) and (l)}

(l+1) $\forall i[i \in \mathbb{N} : P(i) \Rightarrow P(i+1)]$
{induction on (m) and (l+1)}

(l+2) $\forall n[n \in \mathbb{N} : P(n)]$

Basis

induction
hypothesis

Induction step

Inductive definitions

Inductive proof: truth is passed on

Inductive definition: construction is passed on

well defined by induction

Example

The sequence of real numbers $(a_i \mid i \in \mathbb{N})$ is defined inductively by

$$\begin{aligned} a_0 &= 2 \\ a_{i+1} &= 2a_i - 1 \end{aligned}$$

a	a	a	a	a	...
2	3	5	9	17	...

proof by induction

Conjecture

For all $n \in \mathbb{N}$ it holds that

$$a_n = 2^{n+1}$$

Strong induction

P - unary predicate
over \mathbb{N}

$$\forall k [k \in \mathbb{N} : \forall j [j \in \mathbb{N} \wedge j < k : P(j)] \Rightarrow P(k)] \Rightarrow \forall n [n \in \mathbb{N} : P(n)]$$

\forall elim with $k=1$

\Rightarrow elim,
 \wedge intro



$$\begin{aligned} &P(0) \\ &P(0) \Rightarrow P(1) \\ &P(0) \wedge P(1) \\ &P(0) \wedge P(1) \Rightarrow P(2) \\ &P(0) \wedge P(1) \wedge P(2) \\ &P(0) \wedge P(1) \wedge P(2) \Rightarrow P(3) \\ &\dots \end{aligned}$$

Definition of
 $(a_i \mid i \in \mathbb{N})$
with strong
induction

a_n is defined via
 a_0, \dots, a_{n-1}

Cardinality

Cardinals

Def.

Two sets A and B have the same cardinality (are equinumerous) if there is a bijection $f:A \rightarrow B$.
Notation $A \sim B$, or $|A| = |B|$.

Prop.

The relation \sim is an equivalence relation on sets.

Def.

A set A has at most as large cardinality as a set B if there is an injection $f:A \rightarrow B$.
Notation $|A| \leq |B|$.

Def.

A set A has at least as large cardinality as a set B if there is a surjection $f:A \rightarrow B$.
Notation $|A| \geq |B|$.

Def.

A set A has smaller cardinality than a set B if there is an injection $f:A \rightarrow B$ and there is no surjection $f:A \rightarrow B$. Notation $|A| < |B|$.

$$|A| = [A]_{\sim}$$

cardinal
numbers are
 \sim equivalence
classes

Theorem (Cantor)

If $|A| \leq |B|$
and
 $|B| \leq |A|$,
then
 $|A| = |B|$.

Operations on cardinals

Def.

Let A and B be two disjoint sets. Then $|A| + |B| = |A \cup B|$.

Def.

Let A and B be two sets. Then $|A| \cdot |B| = |A \times B|$.

Def.

Let A and B be two sets. Then $|A|^{|B|} = |A^B|$ where A^B is the set of all functions from B to A , i.e. $A^B = \{f \mid f: B \rightarrow A\}$.

Prop.

Let A be a set. Then $|\mathcal{P}(A)| = 2^{|A|}$.

$$|A| = [A]_{\sim}$$

cardinal numbers are \sim equivalence classes

Note: $2 = |\{0, 1\}|$

Finite sets, finite cardinals

We write \mathbb{N}_k for the set $\{0, 1, \dots, k-1\}$. Then $\mathbb{N}_0 = \emptyset$.

We will also write k for $|\mathbb{N}_k|$.

Def.

A set A is finite if and only if $|A| = k$, for some $k \in \mathbb{N}$.

Hence

A set A is finite if and only if there is a natural number $k \in \mathbb{N}$ and a bijection $f: A \rightarrow \mathbb{N}_k$.

$$|A| = [A]_{\sim}$$

cardinal numbers are \sim equivalence classes

if and only if A has k elements, for some $k \in \mathbb{N}$

E.g. If $|A| = k$ and $|B| = m$ for some $k, m \in \mathbb{N}$ then $|A \times B| = k \cdot m$

The operations on cardinals when restricted to finite cardinals coincide with the operations on natural numbers!
This justifies the notation.

Infinite, countable and uncountable sets

Time for a video!

Hilbert's
infinite hotel :-)